

NUTRIPLANT® AG

6-4-3

Low Analysis Fertilizer

A Foliar Nutritional Supplement to Enhance Growth of Agricultural Crops

Guaranteed Analysis:

| | | | | | |
|--|-------|---|------------------------------------|--------|---|
| Total Nitrogen (N) | 6.0 | % | Copper (Cu) | 0.28 | % |
| 3.3% Ammoniacal Nitrogen | | | 0.28% Water Soluble Copper (Cu) | | |
| 2.7% Nitrate Nitrogen | | | Iron (Fe) | 0.36 | % |
| Available Phosphate (P ₂ O ₅) | 4.0 | % | 0.36% Water Soluble Iron (Fe) | | |
| Soluble Potash (K ₂ O) | 3.0 | % | Manganese (Mn) | 0.30 | % |
| Sulfur (S) | 1.0 | % | 0.30% Water Soluble Manganese (Mn) | | |
| 1.0% Combined Sulfur (S) | | | Molybdenum (Mo) | 0.0009 | % |
| Boron (B) | 0.02 | % | Zinc (Zn) | 0.60 | % |
| Cobalt (Co) | 0.015 | % | 0.60% Water Soluble Zinc (Zn) | | |

Derived from Ammonium Nitrate, Ammonium Phosphate, Dipotassium Phosphate, Boric Acid, Cobalt Nitrate, Copper Sulfate, Ferrous Sulfate, Manganese Sulfate, Ammonium Molybdate and Zinc Sulfate.

Chlorine (Cl) not more than 0.5%

F1380

Information concerning the raw materials composing **NUTRIPLANT® AG** can be obtained by writing to Amway Corp. (see address on the label).

**May be Harmful if Swallowed. May Cause Skin and Eye Irritation.
Keep Out of Reach of Children.**

2.5 gal (9.5 liter)

Net weight: 24 lb (11 kg)

Distributed and Guaranteed by

Access Business Group International LLC

7575 Fulton Street East

Ada, MI 49355.

U.S.A.

Satisfaction Guaranteed

60-3480

LACCF52 ADBE 022678

Recommended NUTRIPLANT® AG Application Rates and Times for Selected Crops

APPLICATION: SHAKE WELL before using. Dilute with water to obtain uniform coverage. Dilution depends on spraying equipment. For high volume sprayers dilute 1:200 to 1:1000. For low volume sprayers dilute 1:20 to 1:100. To co-apply with agrochemicals, dilute 1 part NUTRIPLANT® AG with 5 parts water prior to mixing with pre-diluted chemicals. Because available formulations of agrochemicals differ widely, a small test area should be treated prior to large scale mixing to determine that no undesirable effects occur. For best results apply in the evening or early morning.

| CROP | APPLICATION RATES fl oz/acre | APPLICATION TIME(S) |
|--|---------------------------------|---|
| Alfalfa | 16 | Apply at about 90% ground cover. Repeat application after each cutting. |
| Barley, oats, rye, wheat | 16 | 1 application - winter varieties at first spring growth; spring varieties at 2-5 leaf stage. |
| Berries. | 16 | 2 applications - early to 30% bloom and at fruit set. For multiple picking, after each harvest. |
| Citrus | 14 | 3 applications - pre-bloom, fruit set and at fruit color change. |
| Corn, (field, pop, sweet) | 16 | 1 application - 6-8 leaf stage. |
| Cotton | 8 | 4 weekly applications beginning at pinhead square. |
| Flax/linseed | 16 | 1 application - 7-8 leaf stage. |
| Fruit trees. | 16 | 3 applications - leaf out, fruit set, and at fruit color change. |
| Fruiting vegetables. | 12 | 3 applications - pre-bloom, fruit set and at color change. |
| Grapes | 16 | 3 applications - pre-bloom, fruit set and again 2-3 weeks later. |
| Hops | 16 | 1 application - pre-bloom stage. |
| Jute | 16 | 1 application - intense growth stage. |
| Legumes. | 16 | 2 applications - pre-bloom and pod set stage. |
| Melons. | 12 | 3 applications - pre-bloom, fruit set and at fruit color change. |
| Millo, millets, sorghum | 16 | 1 application - 6-8 leaf stage. |
| Nuts | 16 | 3 applications - pre-bloom, fruit set and again 3 weeks later. |
| Olives | 16 | 3 applications - pre-bloom, fruit set and at fruit color change. |
| Onions, chives, garlic, leeks, shallots | 8 | 3 applications - 6-8 leaf, 2 weeks later, and again 4 weeks later. |
| Peanuts | 16 | 2 applications - pre-bloom and at initial pegging. |
| Potatoes, yam | 14 | 2 applications - 6-8 leaf stage and again 4 weeks later. |
| Rapeseed, canola | 16 | 1 application - winter varieties at first spring growth; spring varieties at 6-8 leaf stage. |
| Rice | 14 | 2 applications - 2-5 leaf stage and at panicle initiation. |
| Root vegetables | 8 | 3 applications - on a bimonthly basis, starting at 4-6 leaf stage. |
| Safflower, sesame | 16 | 1 application - pre-bloom stage. |
| Soybeans | 16 | 2 applications - pre-bloom and pod set stage. |
| Strawberries | 8 | At early bloom (up to 30%). For multiple picking, after each harvest. |
| Stem, leaf flower vegetables | 10 | 3 applications - 4-5 leaf stage, 2 weeks later and again 4 weeks later. |
| Sugar beets | 16 | 1 application - 8-10 leaf stage. |
| Sugar cane | 16 | 2 applications - 45-70 days after planting and 6 weeks later. |
| Sunflowers | 16 | 1 application - at star (bud) formation. |
| Tobacco | 16 | 2 applications - 3 weeks after transplanting and again 6 weeks after transplanting. |
| Tropical fruits | 16 | 2 applications - at the beginning of intensive plant growth and at fruit set. |
| Turfigrass* | 16 | 3-5 applications per season, 1-2 months apart (0.37 fl oz/1000 sq ft). |

* This product is not a lawn maintenance fertilizer and at recommended application rates delivers less than 0.002 lb of N and P₂O₅ per 1,000 sq ft.

•Blackberry, blueberry, cranberry, currants, dewberry, gooseberry, loganberry, raspberry.

•Apple, apricot, avocado, cherry, nectarine, peach, pear, plum.

•Cucumber, eggplant, pepper, pumpkin, squash, tomato.

•Beans, (dry, faba, green, etc.) chickpeas (garbanzos), lentils, peas.

•Cantaloupe, crenshaw, honeydew, watermelon.

•Beet, carrot, parsnip, radish, turnip.

•Artichoke, asparagus, broccoli, Brussels sprouts, cabbage, cauliflower, celery, endive, lettuce, parsley, spinach.

•Bananas, mangos, papaya, pineapple.

IMPORTANT NOTICE: Read the entire label. Use only in accordance with the label instructions. Use in conjunction with normal fertilizer program. NUTRIPLANT® AG is intended as a supplement to a regular fertilizer program and will not, by itself, provide all the nutrients normally required by plants.

Information regarding the contents and levels of metals in this product is available on the Internet at <http://www.aapfco.org/metals.htm>

STORAGE: Store at a temperature between freezing and 110°F (43°C). Keep container tightly closed. Do not store diluted product. Do not store in direct sunlight.